


GEOLOGIC MAP OF THE OWINZA BUTTE QUADRANGLE, JEROME AND LINCOLN COUNTIES, IDAHO

John W. Shervais, Scott H. Matthews,
John D. Kauffman, and Kurt L. Othberg
2006

Disclaimer: This Digital Web Map is an informal report and may be revised and formally published at a later time. Its content and format may not conform to agency standards.

CORRELATION OF MAP UNITS



INTRODUCTION

The geologic map of the Owinza Butte quadrangle identifies both the bedrock and surficial geologic units. It shows the geographic distribution of rock types at the surface and in the shallow subsurface. The Owinza Butte quadrangle lies near the center of the Snake River Plain, a large arcuate, lava-filled depression crossing southern Idaho. Pleistocene basalt flows from shield volcanoes, such as Owinza Butte, form the land surface. The older basalt flows are mantled with alluvium and wind-blown sand and silt which form the soils that are cultivated. The geologic units in the area control soil development, groundwater movement and recharge, and geotechnical factors important in construction design and waste management. Land uses in the area include grazing, irrigated agriculture, and dairy farms with confined animal feeding operations. The Snake River Plain aquifer underlies the area and discharges to the southwest of the Owinza Butte quadrangle as springs in the Snake River Canyon.

Modern geologic mapping of the Owinza Butte quadrangle was started through the U.S. Geological Survey EDMAP program, which supported work by Matthews (2000) and Shervais and Matthews (2004). With support from the U.S. Geological Survey's STATEMAP program, additional field investigations by the Idaho Geological Survey of both bedrock and surficial geology completed the mapping. Earlier geologic mapping was by Malde and others (1963). Exposures of the geology were examined in the field and selectively sampled. Matthews (2000) provides results and interpretation of basalt-sample chemical analysis. Aerial photographs were studied to aid in identifying boundaries between map units through photogeologic mapping of landforms. Soil series information is from Ames (2003) and Johnson (2002). The information depicted at this scale furnishes a useful overview of the area's geology but is not a substitute for site-specific evaluations.

DESCRIPTION OF MAP UNITS

ALLUVIAL DEPOSITS

Qas Alluvium of side-streams (Holocene)—Silt and sand flood-plain and sheet-wash deposits in the drainage systems formed between older and younger basalt units.

MIXED LACUSTRINE AND ALLUVIAL DEPOSITS

Qlp Playa deposits (Holocene and Pleistocene)—Fine sand, silt, and clay sorted into thin beds and laminae. Sediments largely derived from erosion of loess from surrounding basalt surfaces and washed into areas of internal drainage or nearly flat slopes. Form flat to gently sloping fills in shallow depressions primarily between basalt flows. Deposited during periodic floods, especially during periods of heavy rains and times of rapid snow melt. These conditions were probably more prevalent during the Pleistocene, therefore the deposits are mostly relict.

BASALT UNITS

The surface geology of the Snake River Plain north of the Snake River is primarily Pleistocene basalt flows of the Snake River Group. On the Owinza Butte quadrangle, the basalt flows originate from several shield volcanoes within and beyond the borders of the quadrangle. Each volcano probably extruded numerous lava flows or flow lobes, although individual flows cannot easily be mapped, especially on the older surfaces now subdued by surficial deposits. Nearly all of the basalt is vesicular to extremely vesicular and most of the units are also diktytaxitic to some degree (i.e., containing voids with protruding crystals). Even units with a fine-grained groundmass have a coarse, grainy texture. Older basalt surfaces tend to be less rugged and more subdued than younger surfaces, primarily the result of greater accumulation of loess over a longer period of time. Over time, drainage patterns change from essentially no drainage on young, very rugged topography, to radial drainage on older buttes. Likewise, young basalt surfaces support little or no agriculture because of the lack of soil, while the older surfaces with thin to thick soil support a wide variety irrigated crops and grazing pastures.

Qrb Basalt of Rocky Butte (Pleistocene)—Fine-grained, dark gray to black, glassy basalt with common to abundant olivine grains 0.5-1 mm in diameter. Common to abundant small plagioclase laths to about 1 mm in length. Remanent magnetic polarity is normal, as determined in the field and through laboratory analysis. Erupted from a shield volcano located less than 2 miles south of the quadrangle, which shows a permanent horizontal-control mark labeled "Rocky" at 4526 feet on the south rim of the vent (sec. 14, T. 8. S., R. 20 E.). Equivalent in part to Sand Springs Basalt of Malde and Powers (1962), Malde and others (1963), Covington (1976), and Covington and Weaver (1990). Thin loess covers the surface of the unit except for pressure ridges, and soil caliche is present but generally thin and weakly developed. Some of the land is cultivatable. Tauxe and others (2004) report an ⁴⁰Ar/³⁹Ar weighted mean plateau age of 0.095 Ma for "Sand Springs" basalt. The location of their sample, on the north rim of the Snake River canyon near Shoshone Falls, is from the unit we map as basalt of Rocky Butte.

Qkim Basalt of Kimama Butte (Pleistocene)—Fine-grained, abundantly plagioclase- and olivine-phyric basalt. Plagioclase phenocrysts 5-10 mm in length. Common to abundant olivine grains 1-2 mm in diameter. Erupted from Kimama Butte, located 6 miles southeast of the quadrangle. Remanent magnetic polarity uncertain. Both normal and reverse polarities were obtained in the field. Duane Champion (written communication, 2004) reports normal polarity based on laboratory analysis of a sample collected from a flow several miles northeast of Kimama Butte.

Qob Basalt of Owinza Butte (Pleistocene)—Medium-grained, black plagioclase- and olivine-phyric basalt. Plagioclase phenocrysts are lath shaped and range from 0.8-4.0 mm in length. Olivine phenocrysts are 0.3-1.3 mm in diameter. Erupted from Owinza Butte. Much of the basalt surface is covered with soil and thin eolian deposits, but some pressure ridges and collapsed lava tubes are still visible.

Qsid Basalt of Sid Butte (Pleistocene)—Basalt flows originating from Sid Butte, located 2 miles east of the quadrangle. Not examined in detail. Cinder pit near the summit contains gabbroic xenoliths (Matthews, 2000). Remanent magnetic polarity is reverse as determined in the laboratory (Duane Champion, written communication, 2004) and in the field during this project.

QTKn Basalt of Knoll 4610 (Pleistocene or Pliocene)—Fine-grained, dark gray basalt with common plagioclase phenocrysts. Remanent magnetic polarity is reverse, as determined in the field. Equivalent to Baby Butte basalt of Matthews (2000) and Shervais and Matthews (2004). Exposures are poor and limited to pavement outcrops on top of the knoll and scattered small outcrops on the flanks. Surface mostly soil covered. The unit is flanked on the north, east, and south by flows from Kimama Butte, and on the west by flows from Rocky Butte.

SYMBOLS

Contact: Line showing the approximate boundary between one map unit and another. The location accuracy of an approximate contact is 80 feet or more on the ground.

Lava flow front: Edge of younger lava flow that erupted onto an older flow from the same source. Includes individual cooling fronts formed during the same eruption.

REFERENCES

Ames, Dal, 2003. Soil survey of Jerome County and Part of Twin Falls County, Idaho: U.S. Department of Agriculture, Natural Resources Conservation Service, 391 pages, 67 sheets.

Covington, H.R., 1976. Geologic map of the Snake River Canyon near Twin Falls, Idaho: U.S. Geological Survey Miscellaneous Field Studies Map MF-809, 2 sheets, scale 1:24,000.

Covington, H.R., and J.N. Weaver, 1990. Geologic map and profiles of the north wall of the Snake River Canyon, Jerome, Filer, Twin Falls, and Kimberly quadrangles, Idaho: U.S. Geological Survey Miscellaneous Investigations Series, Map I-1947-D, scale 1:24,000.

Grunh, Ruth, 1961. Archaeology of Wilson Butte Cave, Idaho Museum of Natural History, Special Paper, Pocatello, ID.

Johnson, M.E., 2002. Soil survey of Wood River area, Idaho, Gooding County and parts of Blaine, Lincoln, and Minidoka counties: U.S. Department of Agriculture, Natural Resources Conservation Service, 797 pages, online at http://ftp-fc.sc.egov.usda.gov/MOI/text_pdf/ida/oid681_text.pdf.


Malde H.E., and H.A. Powers, 1962. Upper Cenozoic stratigraphy of western Snake River Plain, Idaho: Geological Society of America Bulletin, v. 73, p. 1197-1220.

Malde, H.E., Powers, and C.H. Marshall, 1963. Reconnaissance geologic map of west-central Snake River Plain, Idaho: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-373.



Matthews, S.H., 2000. Geology of Owinza Butte, Shoshone SE, and Star Lake quadrangles, Snake River Plain, southern Idaho: University of South Carolina M.S. thesis, 110 p.

Shervais, J.W., and S.M. Matthews, 2004. Geologic map of the Owinza Butte 7.5' quadrangle, Lincoln and Jerome counties, Idaho: EDMAP deliverable, scale 1:24,000.

Tauxe, Lisa, Casey Luskin, Peter Selkin, Phillip Gans, and Andy Calvert, 2004. Paleomagnetic results from the Snake River Plain: contribution to the time-averaged field global database: Geochimica Geophysica Geosystems (G³), v. 5, no. 8, Q08H13 DOI 10.1029/2003GC000661.



Base map from USGS digital raster graphic base, 1992.
Topography by photogrammetric methods from aerial photographs taken 1969. Updated from aerial photographs taken 1987. Field checked 1987. Map edited 1992.
Transverse Mercator, 1927 North American Datum.
10,000-foot grid ticks based on Idaho coordinate system, west zone.
1000-meter Universal Transverse Mercator grid ticks, zone 11.



Field work conducted 2004.
This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program.
Digital cartography by Jane S. Freed at the Idaho Geological Survey's Digital Mapping Lab.
Map version 8-30-2006.
Note on printing: The map is reproduced at a high resolution of 600 dots per inch. The inks are resistant to run and fading but will deteriorate with long-term exposure to light.
PDF map (Acrobat Reader) may be viewed at www.idahogeology.org.